arXiv:1304.2982v1 [cond-mat.stat-mech] 10 Apr 2013

Non-mean-field Critical Exponent in a Mean-field Model : Dynamics versus Statistical Mechanics

Shun Ogawa,[1] Aurelio Patelli,[2] and Yoshiyuki Y. Yamaguchi[1]
[1]Department of Applied Mathematics and Physics, Graduate School of Informatics, Kyoto University, Kyoto, 606-8501, Japan
[2]Dipartimento di Fisica ed Astronomia, Universitá di Firenze and INFN, Via Sansone 1, IT-50019 Sesto Fiorentino, Italy

(Dated: April 11, 2013)

The critical exponent of susceptibility is expected as 1 in systems belonging to the mean-field universality class. On the other hand, the mean-field nature permits to introduce the Vlasov equation in the place of the Liouville equation, and the linear response theory based on the Vlasov equation gives the strange exponent $1/4$ in the low-temperature side of the Hamiltonian mean-field model. We clarify that this strange exponent is due to existence of Casimir invariants which traps the system in a quasistationary state for a time scale diverging with the system size. The theoretical prediction is numerically confirmed by N-body simulations and numerical Vlasov solutions.

PACS numbers: 05.20.Dd, 64.60.fh, 75.40.Cx

Is the universality of critical exponents extended from the context of statistical mechanics to of dynamics? For a ferromagnetic model, we can define the isothermal susceptibility χ^T and the static isolated susceptibility $\chi^{\rm I}$. The former comes from the statistical mechanics of isothermal processes, and the latter comes from dynamical arguments and is defined by the linear response theory [1]. These two susceptibilities satisfy the inequality $\chi^T \geq \chi^{\rm I}$ [2, 3] which takes care of the existence of invariants in the considering processes [4, 5]. The above inequality implies a similar inequality between the critical exponents $\gamma_\pm$, since susceptibility is scaled as $\chi \sim |\tau|^{-\gamma_\pm}$ around the critical temperature $T_{\rm c}$. Here $\tau = (T - T_{\rm c})/T_{\rm c}$ is the reduced temperature, T is temperature, and γ_+ and γ_- are defined in the high- and the low-temperature sides respectively. Breaking of the equality may happen between the value of the susceptibilities, that is $\chi^T > \chi^{\rm I}$, but this breaking does not imply the breaking of equality between the corresponding critical exponents.

To tackle this problem, we concentrate on systems in the mean-field universality class, whose critical exponents are $\gamma_\pm = 1$. One of such systems is the Hamiltonian mean-field (HMF) model [6–8], in which the mean-field approximation is exact in the limit of large number of particles. HMF is an important toy model in the studies of long-range interacting systems and, moreover, it is used in the theoretical description of collective atomic recoil laser experiments [9]. The potential term of the HMF model is the same of the globally-coupled XY model, and hence the HMF model is a ferromagnetic model belonging to the mean-filed universality class. Magnetization M in the dynamical context, namely time average, agrees with one in the context of statistical mechanics when initial conditions for N particles are sampled from the canonical equilibrium distribution and the external magnetic field is not introduced [10]. In other words, the initial question is positively answered in the case of the magnetization, which scales as $M \sim (-\tau)^\beta$ with $\beta = 1/2$. This agreement allows to consider the initial question for $\gamma_\pm$, since, to derive the exponents, we have to prepare canonical equilibrium states before adding the external magnetic field.

For the HMF model, we can compute χ^T explicitly and its critical exponents are $\gamma_\pm^T = 1$ [11]. In contrast, the computation of $\chi^{\rm I}$ is not an easy task since we have to solve the N-body dynamics. Nevertheless, the mean-field nature of the HMF model permits to compute a dynamical susceptibility, since it naturally introduces the Vlasov equation in the place of the Liouville equation. Thanks to integrability of the one-particle Hamiltonian associated with the Vlasov equation, the linear response theory based on the Vlasov equation gives the exact susceptibility in the mean-field limit. We call it the Vlasov susceptibility and denote with $\chi^{\rm V}$.

The critical exponent of $\chi^{\rm V}$ is $\gamma_+^{\rm V} = 1$ in the high-temperature homogeneous side [12], but seems smaller than 1 in the low-temperature inhomogeneous side [13]. We will show that $\gamma_-^{\rm V} = 1/4$ due to existence of invariants, called Casimirs. We clarify that existence of the Casimirs traps the system in a quasistationary state (QSS) and induces the strange exponent. We also show that the life time of these QSSs is $O(N)$, and hence, the accessible critical exponent of the susceptibility is not 1 but $1/4$ if the number of particles is large enough.

We remark on studies of the critical exponents in the Vlasov dynamical systems. Based on the theory constructing an unstable manifold in the Vlasov-Poisson equation [14], Ivanov et al. [15] have stated that scaling laws are different from ones predicted by the ordinary mean-field theory. However, they start from an unstable spatially homogeneous Maxwell distribution, and have said nothing about the critical exponents starting from the canonical equilibrium states. We stress that a special choice of initial states may give non-mean-field critical exponents, but our analysis is referred to canonical equilibrium states.

The HMF model consists of N particles confined in the unit circle on the (x, y) plane, and is expressed by the Hamiltonian

$$H_N = \sum_{i=1}^{N} \left[\frac{p_i^2}{2} + \sum_{j=1}^{N} \frac{1 - \cos(q_i - q_j)}{2N} - h\Theta(t)\cos(q_i - \phi) \right] \tag{1}$$

where h and ϕ are the modulus and the phase of the external magnetic vector $(h\cos\phi, h\sin\phi)$ respectively, and $\Theta(t)$

is the Heaviside step function. The magnetization vector $(\langle M_x\rangle_N, \langle M_y\rangle_N)$ is defined by $\langle M_x\rangle_N + i\langle M_y\rangle_N = \sum_{j=1}^{N} e^{iq_j}/N$, where $\langle\cdot\rangle_N$ represents the average over N particles. This system has a rotational symmetry with $h = 0$, and we may set $\langle M_y\rangle_N = 0$ without loss of generality. We assume $\phi = 0$ to concentrate our interests on susceptibility along the direction of the spontaneous magnetization, defined x. The magnetization observable is $M = \cos q$, and its average over the one-particle distribution $f(q,p,t)$ is denoted by $\langle M\rangle = \iint M f dq dp$. The corresponding effective one-particle Hamiltonian is written as

$$H[f](q,p,t) = p^2/2 - \langle M\rangle \cos q - h\Theta(t)\cos q, \tag{2}$$

where we denote $\langle M\rangle = \iint M f dq dp$ the average over the one-particle distribution function $f(q,p,t)$. This distribution is a solution to the Vlasov equation

$$\partial_t f + \partial_p H[f]\partial_q f - \partial_q H[f]\partial_p f = 0, \tag{3}$$

associated with the effective Hamiltonian (2).

To clarify the discrepancy between the isothermal susceptibility and the Vlasov susceptibility, we first compute the isothermal susceptibility for the canonical equilibrium state with temperature T. The mean-field approximation, which is exact in the limit $N \to \infty$ in the HMF model, allows to compute magnetization by the one-particle distribution function

$$f_h = e^{-[p^2/2-(\langle M\rangle_h+h)\cos q]/T} \Bigg/ \iint e^{-[p^2/2-(\langle M\rangle_h+h)\cos q]/T} dq dp\ , \tag{4}$$

where $\langle M\rangle_h$ is the average of M with respect to f_h. We put $h = 0$ for $t < 0$, and $h > 0$ for $t > 0$. The equilibrium magnetization $\langle M\rangle_h$ must satisfy the self-consistent equation

$$\langle M\rangle_h = I_1\left((\langle M\rangle_h + h)/T\right) / I_0\left((\langle M\rangle_h + h)/T\right), \tag{5}$$

where I_n is the modified Bessel function of the first kind.

From the definition of the isothermal susceptibility χ^T, the magnetization $\langle M\rangle_h$ is written as

$$\langle M\rangle_h = \langle M\rangle_0 + h\chi^T + O(h^2), \tag{6}$$

and hence f_h is expanded as

$$f_h = \left[1 + h\frac{\chi^T + 1}{T}(M - \langle M\rangle_0)\right] f_0 + O(h^2). \tag{7}$$

Multiplying f_h by M and integrating over q and p, and comparing the obtained expression with (6), the isothermal susceptibility is written in the form

$$\chi^T = (1 - D^T)/D^T \tag{8}$$

where

$$D^T = 1 - \frac{\langle M^2\rangle_0 - \langle M\rangle_0^2}{T}. \tag{9}$$

On the other hand, the Vlasov susceptibility χ^{V} is obtained by introducing the angle-action variables (θ, J) in the Hamiltonian system (2) with $h = 0$ [16]. Thanks to the Vlasov linear response theory [13], the Vlasov susceptibility is given in a similar form

$$\chi^{\mathrm{V}} = (1 - D^{\mathrm{V}})/D^{\mathrm{V}}, \tag{10}$$

and

$$D^{\mathrm{V}} = 1 - \frac{\langle M^2\rangle_0 - \langle C^2\rangle_0}{T}, \tag{11}$$

where C is the average of $\cos q$ over the angle θ,

$$C(J) = \frac{1}{2\pi}\int_0^{2\pi} \cos q(\theta, J) d\theta, \tag{12}$$

and $\langle C^2\rangle_0 = \iint C^2 f_0 d\theta dJ$. We compare the scaling of the Vlasov susceptibility (10) with the one of the isothermal susceptibility (8), both in the high- and the low-temperature sides.

In the high-temperature side we have $\langle M\rangle_0 = 0$ and $C = 0$, since the angle variable is q itself, and $\langle M^2\rangle_0 = 1/2 = T_{\mathrm{c}}$. We therefore have $D^T = D^{\mathrm{V}} = (T - T_{\mathrm{c}})/T$ and

$$\chi^T = \chi^{\mathrm{V}} = \tau^{-1}, \tag{13}$$

which implies that the critical exponents are the same and of the mean-field universality class: $\gamma_+^T = \gamma_+^{\mathrm{V}} = 1$. The theoretical prediction for χ^{V} is confirmed by direct numerical simulations of the Vlasov equation for small enough external field h as shown in Fig. 1. The discrepancy between the canonical prediction and the numerics in Fig. 1 comes from nonlinear effects. No QSS features can be seen in a small h region where the linear response theory is valid.

In the low-temperature side, we separately analyze the common term $1 - \langle M^2\rangle_0/T$ and the individual terms between D^T and D^{V}. The common part approaches to zero linearly again in the limit $\tau \nearrow 0$. Indeed, the mean square of M is $\langle M^2\rangle_0 = 1 - T$. Thus, we find $1 - \langle M^2\rangle_0/T \sim 2\tau$. Moreover, the individual term of the isothermal function D^T (9) shows the scaling $\langle M\rangle_0^2 \sim (-\tau)^{2\beta}$ with $\beta = 1/2$ of the mean-field universality class. The isothermal susceptibility, therefore, shows the classical exponent $\gamma_-^T = 1$ as expected by the statistical mechanics of mean-field models. On the Vlasov counterpart, by using the new variable $k = \sqrt{(H[f_0] + \langle M\rangle_0)/(2\langle M\rangle_0)}$, the individual term $\langle C\rangle_0^2$ in D^{V} (11) can be expressed as [13]

$$\begin{aligned}\langle C^2\rangle_0 &= 16\sqrt{\langle M\rangle_0}\int_0^1 \left[\frac{2E(k)}{K(k)} - 1\right]^2 kK(k) f_0(k) dk \\ &\quad + 16\sqrt{\langle M\rangle_0}\int_1^\infty \left[\frac{2k^2E(1/k)}{K(1/k)} - 2k^2 + 1\right]^2 K(1/k) f_0(k) dk,\end{aligned} \tag{14}$$

where $f_0(k) = A_0 \exp[\langle M\rangle_0(1 - 2k^2)/T]$ and A_0 is the normalization factor. We note that the two integrals of (14) are positive and finite. In the limit of $k \nearrow 1$, the divergence of



FIG. 1: (color online) Short time average of magnetization, $\overline{M}$, divided by h in the high-temperature side. $T = 0.51$ (red circle), 0.55 (green triangle) and 0.60 (blue diamond). Averages are taken in the period $100 < t \leq 3000$. Temporal evolutions are obtained by the semi-Lagrangian method [17], and the grid sizes are 256×512 (filled symbols) and 128×256 (empty symbols) in $(q, p) \in (-\pi, \pi] \times [-5, 5]$. Time slice is 0.05. Horizontal lines mark theoretical predictions of susceptibility (13) for $T = 0.51$, 0.55 and 0.60 from top to bottom. The broken lines are the predictions given by the canonical ensemble for the same set of temperature.



FIG. 2: (color online) The Vlasov susceptibility $\chi^{\rm V}$ (red solid lower) and the isothermal susceptibility χ^T (green broken upper) as functions of $-\tau$ in log-log plot. The adiabatic susceptibility χ^S (blue dashed middle) is also reported, which is computed explicitly by using the exact solution in the microcanonical statistics [8] and by taking the invariance of the entropy during the quasistatic adiabatic process into account. The black straight lines are guides of eyes for the slopes $-1/4$ and -1. Purple square points are computed by time averages over the period $0 \leq t \leq 500$ in N-body simulations with $N = 10^6$ and time slice 0.1. For each T, 10 points are plotted corresponding to 10 realizations.

$K(k)$ is written as $\log(1 - k^2)$ and is integrable. In a large k region of the second integral, damping of $f_0(k)$ becomes slower as $T \nearrow T_{\rm c}$ since $\langle M \rangle_0 \searrow 0$. However, the factor of the brackets gives $O(k^{-4})$ damping, then the second integral converges. Consequently, $\langle C^2 \rangle_0$ approaches to zero in the limit $\tau \nearrow 0$ as $\langle C^2 \rangle_0 \sim \sqrt{\langle M \rangle_0} \sim (-\tau)^{\beta/2}$, where we used the agreement of $\langle M \rangle_0$ between the contexts of statistical mechanics and of dynamics [10]. The function $D^{\rm V}$ is hence written in the form $D^{\rm V} = c_1|\tau| + c_2|\tau|^{\beta/2}$, and the second term dominates around $\tau \lesssim 0$. Accordingly, the susceptibility $\chi^{\rm V}$ scales as $\chi^{\rm V} \sim (-\tau)^{-\beta/2}$ and the Vlasov critical exponent is $\gamma_-^{\rm V} = \beta/2 = 1/4$. This exponent is numerically confirmed by computing the two integrals in (14) and by N-body simulations as shown in Fig. 2.

The strange divergence of the Vlasov susceptibility comes from the term $\langle C^2 \rangle_0$, and we will show that this term comes from the existence of Casimir invariants. A Casimir is written by using a smooth function s as $\int s(f(q,p,t))dqdp$ and is an integral of the Vlasov equation even if the Hamiltonian $H[f]$ depends on time. The distribution function with the external field must keep the Casimir invariants, thus it is no more f_h but $f_{\rm c}$. Writing $f_{\rm c} = f_0 + g$ we have the restriction for g so that every Casimirs are invariant up to the linear order. We remark that f_0 depends on (q,p) solely through $H[f_0]$, and accordingly, is a function of the action J only. Using the Fourier transform of g with respect to the angle θ, the above restriction is rewritten as

$$0 = \iint [s(f_0 + g) - s(f_0)]dqdp = \int s'(f_0(J))\,\tilde{g}_0(J)dJ \quad (15)$$

where $\tilde{g}_0(J)$ is the Fourier zero-mode of g. This constraint must hold for any smooth functions s, and therefore, we conclude that $\tilde{g}_0(J) = 0$ [18]. We remark that the normalization condition for $f_{\rm c}$ automatically holds accordingly.

The above observation constructs $g = f_{\rm c} - f_0$ by subtracting the Fourier zero-mode from $f_h - f_0$. For this purpose, we expand f_h as (7) but replace χ^T with $\chi^{\rm V}$. The Fourier zero-mode of $f_h - f_0$ is expressed by

$$\frac{1}{2\pi}\int_0^{2\pi}(f_h - f_0)d\theta = h\frac{\chi^{\rm V}+1}{T}(C(J) - \langle M \rangle_0)f_0 + O(h^2). \quad (16)$$

Consequently, the constructing distribution function $f_{\rm c}$ is

$$f_{\rm c} = f_0 + h\frac{\chi^{\rm V}+1}{T}(M - C(J))f_0 + O(h^2). \quad (17)$$

The distribution function $f_{\rm c}$ gives magnetization $\langle M \rangle_{\rm c} = \iint M f_{\rm c} dqdp$ in the form

$$\langle M \rangle_{\rm c} = \langle M \rangle_0 + h\frac{\chi^{\rm V}+1}{T}(\langle M^2 \rangle_0 - \langle C^2 \rangle_0) + O(h^2). \quad (18)$$

This equation and the expansion $\langle M \rangle_{\rm c} = \langle M \rangle_0 + h\chi^{\rm V} + O(h^2)$ gives the Vlasov susceptibility $\chi^{\rm V}$ (10) and the function $D^{\rm V}$ (11).

Following the above discussions, we propose a scenario of relaxation as follows [19, 20]. For $t > 0$, the system is trapped in a QSS to keep the Casimir invariants and this gives the strange critical exponent $\gamma_-^{\rm V} = 1/4$. However, the Vlasov dynamics is not the true dynamics for finite systems, thus the Casimirs are not exactly invariants but evolve in a time scale which diverges with N. Consequently, the system goes to the thermal equilibrium recovering the classical exponent after a long time.

The above scenario is examined by direct N-body simulations in Fig. 3. For $t < 0$, the system is in thermal equilibrium



FIG. 3: (color online) N-body simulations of the HMF model with external field. $N = 10^3$ (100), 10^4 (10) and 10^5 (1), where the inside of braces are the number of realizations over which the orbits are averaged. The system is in thermal equilibrium with $T = 0.499$ in $t < 0$, and the external field turns on with $h = 0.01$ at $t = 0$. The direction of magnetization vector (M_x, M_y) is reset to x-direction at $t = 0$. In each panel, three horizontal lines represent the thermal equilibrium level without external field (lower), the QSS level predicted by the Vlasov linear response theory (middle) and the thermal equilibrium level with the external field (upper). (a) Short time evolution of magnetization. (b) Long time evolution. The horizontal axis is in the logarithmic scale, which is scaled as $\log_{10}(t/N)$ in the inset.

with the temperature $T = 0.499 < 1/2$. The external field with $h = 0.01$ is switched on at $t = 0$, and the system jumps to the QSS predicted by the linear response theory based on the Vlasov equation. In the long time regime, the Casimirs are no more invariant due to appearance of the collision term, and the system goes towards thermal equilibrium. Simulations indicate that the time scale of relaxation from the QSS to thermal equilibrium grows linearly with N, as found in inhomogeneous QSSs without external fields [21].

Finally five remarks are in order.

The first remark is on the validity of the linear response theory to discuss critical behaviours. The linear response theory assumes that $f_c - f_0$ is small enough, and it seems not possible to discuss divergence of susceptibility at the first sight. However, for instance in the low-temperature side, what we have to keep is $|h\chi^{\rm V}| \ll \langle M\rangle_0$. Thus, the linear response theory may be valid even around the critical point by setting the external field to be small enough. The Vlasov susceptibility is computed by taking the limit of $h \to 0$, and hence the Vlasov susceptibility can be computed by the Vlasov linear response theory. See Fig. 1 for the high-temperature side.

The second remark is on the universality of the exponent $\gamma_-^{\rm V} = \beta/2$. We can estimate $D^{\rm V}$ for any QSSs by using the formula shown in [13]. We obtain $D^{\rm V} \sim \sqrt{\langle M\rangle_0}$ around the critical points for any families of distribution functions f_0 which depend on (q, p) only through the effective one-particle Hamiltonian $H[f_0]$ and are stable stationary solutions to the Vlasov equation. Thus, the exponent $\gamma_-^{\rm V} = \beta/2$ is obtained for such a family of distribution functions.

The third remark is on the spectrum method to compute susceptibility [22]. The method gives the correct susceptibility in the high-temperature side, which is consistent with the present result, but does not in the low-temperature side, since the existence of integrals is not fully included. The spectrum method could be improved by considering the Casimir integrals.

The fourth remark is that the existence of invariants may break some thermodynamic laws. Indeed, local temperature in isolated crystalline clusters is not uniform by conservation of angular and translational momenta [23].

We end this article by remarking on the difference between dynamics and statistical mechanics. The dynamical system gets trapped in QSSs for a long time, therefore, measures on experimental setups will show the Vlasov prediction for systems with large enough number of particles.

The authors thank to S. Ruffo and C. Nardini for useful discussions and to M. M. Sano and M. Suzuki for informing about Ref. [5]. This work is inspired by discussions in the workshop held in Centre Blaise Pascal, ENS-Lyon, August 2012. YYY acknowledges the support of Grant-in-Aid for Scientific Research (C) 23560069. SO acknowledges the support of Grants for Excellent Graduate Schools and the hospitality of University of Florence.

---

[1] R. Kubo, J. Phys. Soc. Jpn. **12**, 570 (1957).
[2] H. Falk, Phys. Rev. **165**, 602 (1968).
[3] R. M. Wilcox, Phys. Rev. **174**, 624 (1968).
[4] P. Mazur, Physica **43**, 533 (1969).
[5] M. Suzuki, Physica **51**, 277 (1971).
[6] S. Inagaki and T. Konishi, Publ. Astron. Soc. Jpn. **45**, 733 (1993).
[7] M. Antoni and S. Ruffo, Phys. Rev. E **52**, 2361 (1995).
[8] A. Campa, et al, Phys. Rep. **480**, 57 (2009).
[9] R. Bonifacio and L. de Salvo Souza, Nucl. Instrum. Meth. Phys. Res. A **341**, 360 (1994).
[10] V. Latora, et al. Phys. Rev. Lett. **80**, 692 (1998); Physica D **131**, 38 (1999).
[11] P. H. Chavanis, Eur. Phys. J. B **80**, 275 (2011).
[12] A. Patelli et al., Phys. Rev. E **85**, 021133 (2012).
[13] S. Ogawa and Y. Y. Yamaguchi, Phys. Rev. E **85**, 061115 (2012).
[14] J. D. Crawford, Phys. Plasmas **2**, 97 (1995).
[15] A. V. Ivanov et al., Phys. Rev. E **71**, 056406 (2005).

[16] J. Barré et al., J. Stat. Mech. P08002 (2010).
[17] P. de Buyl, Commun. Nonlin. Sci. Num. Sim. **15**, 2133 (2010).
[18] S. Ogawa, arXiv:1301.1130.
[19] Y. Y. Yamaguchi et al., Physica A **337**, 36 (2004).
[20] J. Barré et al., Physica A **365**, 177 (2006).
[21] P. de Buyl et al., Phys. Rev. E **84**, 061151 (2011).
[22] A. Patelli and S. Ruffo, in preparation.
[23] T. Niiyama et al., Phys. Rev. Lett. **99**, 014102 (2007).